

Off-critical Luttinger Junctions

Brando Bellazzini¹, Mihail Mintchev² and Paul Sorba³

¹ *Institute for High Energy Phenomenology Newman Laboratory of Elementary Particle Physics, Cornell University, Ithaca, NY 14853, USA*

² *Istituto Nazionale di Fisica Nucleare and Dipartimento di Fisica dell'Università di Pisa, Largo Pontecorvo 3, 56127 Pisa, Italy*

³ *Laboratoire de Physique Théorique d'Annecy-le-Vieux, UMR5108, Université de Savoie, CNRS, 9, Chemin de Bellevue, BP 110, F-74941 Annecy-le-Vieux Cedex, France*
(Dated: February 1, 2010)

We investigate Luttinger junctions of quantum wires away from criticality. The one-body scattering matrix, corresponding to the off-critical boundary conditions at the junction, admits in general antibound and/or bound states. Their contribution to the theory is fixed by causality. The presence/absence of bound states determines the existence of two different regimes with inequivalent physical properties. A scattering matrix without bound states defines an isolated equilibrium system. Bound states instead drive the system away from equilibrium, giving raise to non-trivial incoming or outgoing energy flows in the junction. We derive in both regimes and in explicit form the electromagnetic conductance tensor, pointing out the different impact of bound and antibound states.

I. INTRODUCTION

The past two decades have shown a constantly growing interest^{1–32} in the physics of quantum wire junctions. The quantum nature of the transport properties of these devices is fairly well described by the Tomonaga-Luttinger model³³ on graphs of the type shown in FIG. 1. The edges $\{E_i : i = 1, \dots, n\}$ of such a *star* graph Γ are modeling the wires, whereas the vertex represents the junction, which can be treated¹⁹ as a point-like defect (impurity) in the Luttinger liquid. The defect is implemented by a nontrivial one-body scattering matrix $\mathbb{S}(k)$, k being the momentum. At criticality the boundary conditions at the vertex are scale invariant and the scattering matrix is a constant $n \times n$ matrix $\mathbb{S}$. Various authors^{10,18,20,23,32} have investigated in this regime the physical properties of the system. In particular, they derived the conductance tensor $\mathbb{G}$, getting the following simple expression

$$\mathbb{G}_{ij} = \mathbb{G}_{\text{line}}(\delta_{ij} - \mathbb{S}_{ij}), \quad (1)$$




FIG. 1: A star graph Γ with scattering matrix $\mathbb{S}(k)$ at the vertex.

the coefficient $\mathbb{G}_{\text{line}}$ being the conductance of a single wire without junctions, which depends on the parameters (see eq. (43) below) of the Luttinger liquid.

In the present paper we pursue further the study of off-critical Luttinger junctions started in Ref. 22. Our main goal here is to extend the analysis to the case when $\mathbb{S}(k)$ admits bound states and to derive the off-critical generalization (see eq. (44) below) of the formula (1). This is not a purely theoretical problem, because realistic quantum wire junctions are usually noncritical. In order to reproduce this situation as close as possible, in what follows we will keep the theory critical in the *bulk* of Γ , allowing for breaking of scale invariance only at the vertex. In this regime the scattering matrix is no longer constant, but depends on the momentum k . The continuation of $\mathbb{S}(k)$ in the complex k -plane is a meromorphic function with poles of the type $k = i\eta$ with real $\eta \neq 0$. It turns out that the behavior of the theory depends in a crucial way on the sign of η . In absence of bound states (all $\eta < 0$), the model has²² a unitary time evolution respecting time-translation invariance. Accordingly, the energy is conserved. The situation radically changes when $\mathbb{S}(k)$ admits bound states (some $\eta > 0$). Each of them generates in the spectrum of the theory a kind of damped harmonic oscillator. These oscillators lead to a breakdown of time-translation invariance. The energy of the system is no longer conserved, which signals a non-trivial energy flow through the boundary (the vertex of Γ). The essential point here is that the relative non-equilibrium state is fixed by the fundamental physical principle of causality (local commutativity). Physically, the off-critical boundary conditions with $\eta > 0$ generate a specific boundary interaction with the environment. One possibility to implement such interaction in realistic quantum wire junctions might be^{9,10,15} an external magnetic field crossing the junction.

The paper is organized as follows. In the next section

we briefly review the Tomonaga-Luttinger (TL) model on a star graph with off-critical boundary conditions at the vertex. The associated scattering matrix $\mathbb{S}(k)$ and its analytic properties are described in detail. The symmetry content of the model is also analyzed here. In section III we examine the impact of the analytic structure of $\mathbb{S}(k)$ on the physical properties of the theory. The generalization of the conductance formula (1) away from criticality is derived as well. The basic features of the off-critical conductance are discussed in detail, confirming the different role played by the bound and the antibound states of $\mathbb{S}(k)$. Section IV is devoted to our conclusions. The Appendix A contains some technical details concerning the quantization on Γ with off-critical boundary conditions.

II. LUTTINGER LIQUID WITH OFF-CRITICAL BOUNDARY CONDITIONS ON Γ

A. Bulk theory and boundary conditions

The dynamics of the Luttinger liquid in the bulk is defined by Lagrangian density

$$\mathcal{L} = i\psi_1^*(\partial_t - v_F\partial_x)\psi_1 + i\psi_2^*(\partial_t + v_F\partial_x)\psi_2 - g_+(\psi_1^*\psi_1 + \psi_2^*\psi_2)^2 - g_-(\psi_1^*\psi_1 - \psi_2^*\psi_2)^2. \quad (2)$$

Here $\{\psi_\alpha(t, x, i) : \alpha = 1, 2\}$ are complex fields, depending on the time t and the position (x, i) , where $x > 0$ is the distance from the vertex and $i = 1, \dots, n$ labels the edge, as shown in FIG. 1. Finally, v_F is the Fermi velocity and $g_\pm \in \mathbb{R}$ are the coupling constants³⁴. The equations of motion following from (2) imply the conservation law

$$\partial_t \rho(t, x, i) - v_F \partial_x j(t, x, i) = 0, \quad (3)$$

where

$$\rho(t, x, i) = (\psi_1^*\psi_1 + \psi_2^*\psi_2)(t, x, i), \quad (4)$$

$$j(t, x, i) = (\psi_1^*\psi_1 - \psi_2^*\psi_2)(t, x, i), \quad (5)$$

are the charge density and electric current respectively. Our main task below is to derive the relative conductance. For this purpose we have to complete first the description of the dynamics, specifying the interaction at the vertex of Γ . In other words, we must fix the boundary conditions on ψ_α at $x = 0$. This is a very delicate point because it strongly interferes with the solution of the model. It is useful to recall in this respect that the model (2) is exactly solvable on the line³³ via bosonization. In order to preserve this nice feature on Γ , it is more convenient to formulate the boundary conditions directly in bosonic terms. We will show now that this strategy, which works^{20–22} nicely at criticality, can be extended to off-critical boundary conditions as well.

The basic ingredient for solving the model (2) via bosonization is the scalar field φ satisfying

$$(\partial_t^2 - \partial_x^2) \varphi(t, x, i) = 0, \quad x > 0 \quad (6)$$

and some boundary conditions for $x = 0$. Following a standard QFT procedure, the latter are fixed by requiring that the operator $K \equiv -\partial_x^2$ on Γ is *self-adjoint*. This problem has been intensively investigated in the recent mathematical literature^{35–37}, where the subject goes under the name of “quantum graphs”. From these studies one infers that K is self-adjoint on Γ if and only if the field φ satisfies the boundary condition^{35,37}

$$\sum_{j=1}^n [\lambda(\mathbb{I} - \mathbb{U})_{ij} \varphi(t, 0, j) - i(\mathbb{I} + \mathbb{U})_{ij} (\partial_x \varphi)(t, 0, j)] = 0, \quad (7)$$

where U is any unitary matrix and $\lambda > 0$ is a parameter with dimension of mass, characterizing the breaking of scale invariance.

Eq. (7) generalizes to the graph Γ the familiar mixed (Robin) boundary condition on the half-line $\mathbb{R}_+$. The matrices $\mathbb{U} = \mathbb{I}$ and $\mathbb{U} = -\mathbb{I}$ define the Neumann and Dirichlet boundary conditions respectively. The physical interpretation of (7) in the context of bosonization was discussed in Ref. 21. At criticality (7) describes the splitting of the electric current (5) at the junction.³⁸

B. Scattering matrix and solution of the model

As already mentioned in the introduction, for the explicit construction of φ it is convenient to interpret¹⁹ the vertex of Γ as a point-like impurity (defect)^{39–42}, characterized by a non-trivial scattering matrix $\mathbb{S}(k)$. This matrix is associated^{35,36} to the operator K on Γ and is fully determined by the boundary conditions (7). The explicit form is³⁵

$$\mathbb{S}(k) = -\frac{[\lambda(\mathbb{I} - \mathbb{U}) - k(\mathbb{I} + \mathbb{U})]}{[\lambda(\mathbb{I} - \mathbb{U}) + k(\mathbb{I} + \mathbb{U})]} \quad (8)$$

and has a simple physical interpretation: the diagonal element $\mathbb{S}_{ii}(k)$ represents the reflection amplitude from the defect on the edge E_i , whereas $\mathbb{S}_{ij}(k)$ with $i \neq j$ equals the transmission amplitude from E_i to E_j .

The matrix (8) has a number of remarkable features. It is unitary $\mathbb{S}(k)^* = \mathbb{S}(k)^{-1}$ by construction and satisfies $\mathbb{S}(k)^* = \mathbb{S}(-k)$, known as Hermitian analyticity. Notice also that $\mathbb{S}(\lambda) = \mathbb{U}$, showing that the boundary condition (7) is fixed actually by the value of scattering matrix at the scale λ .

Let us summarize now the analytic properties of $\mathbb{S}(k)$ needed in what follows. We denote by $\mathcal{U}$ the unitary matrix diagonalizing $\mathbb{U}$ and parametrize

$$\mathbb{U}_d = \mathcal{U}^{-1} \mathbb{U} \mathcal{U} \quad (9)$$

as follows

$$\mathbb{U}_d = \text{diag}(e^{2i\alpha_1}, e^{2i\alpha_2}, \dots, e^{2i\alpha_n}), \quad \alpha_i \in \mathbb{R}. \quad (10)$$

Using (8), one easily verifies that $\mathcal{U}$ diagonalizes also $\mathbb{S}(k)$

for any k and that

$$\mathbb{S}_d(k) = \mathcal{U}^{-1} \mathbb{S}(k) \mathcal{U} = \text{diag} \left(\frac{k + i\eta_1}{k - i\eta_1}, \frac{k + i\eta_2}{k - i\eta_2}, \dots, \frac{k + i\eta_n}{k - i\eta_n} \right), \quad (11)$$

where

$$\eta_i = \lambda \tan(\alpha_i), \quad -\frac{\pi}{2} \leq \alpha_i \leq \frac{\pi}{2}. \quad (12)$$

We conclude therefore that $\mathbb{S}(k)$ is a meromorphic function with *simple* poles located on the imaginary axis and different from 0. In what follows we denote by $\mathcal{P} = \{i\eta : \eta \neq 0\}$ the set of *distinct* poles of $\mathbb{S}(k)$: the subset $\mathcal{P}_+ = \{i\eta : \eta > 0\}$ in the upper half-plane corresponds to *bound* states whereas $\mathcal{P}_- = \{i\eta : \eta < 0\}$ in the lower half-plane gives rise to *antibound* states.⁴³ We will make use below also of the *residue* matrix defined by

$$R_{ij}^{(\eta)} = \frac{1}{i\eta} \lim_{k \rightarrow i\eta} (k - i\eta) \mathbb{S}_{ij}(k), \quad i\eta \in \mathcal{P}. \quad (13)$$

We recall that in the above parametrization the angular variables α_i characterize the departure from criticality, the critical points corresponding²² to the values $\alpha_i = 0, \pm\pi/2$. The classification and the stability properties of these critical points have been extensively studied in the literature^{3,7,10,18,20–23,31}. As already stated in the introduction, our goal here is to go beyond and explore the theory away from criticality.

The basic steps in the construction of the field φ , which satisfies the wave equation (6) and the off-critical boundary condition (7), are given in Appendix A. The subtle point is the contribution of the bound states of $\mathbb{S}(k)$, which need a separate treatment where local commutativity turns out to be essential. The dual field $\tilde{\varphi}$ is defined by

$$\partial_t \tilde{\varphi}(t, x, i) = -\partial_x \varphi(t, x, i), \quad (14)$$

$$\partial_x \tilde{\varphi}(t, x, i) = -\partial_t \varphi(t, x, i), \quad (15)$$

and the solution of the Luttinger model on Γ can be expressed in terms of the pair $\{\varphi, \tilde{\varphi}\}$. For the details we refer to Refs. 20–22, recalling here only the solution

$$\psi_1(t, x, i) \sim : e^{i\sqrt{\pi}[\zeta_+ \varphi(vt, x, i) - \zeta_- \tilde{\varphi}(vt, x, i)]} :, \quad (16)$$

$$\psi_2(t, x, i) \sim : e^{i\sqrt{\pi}[\zeta_+ \varphi(vt, x, i) + \zeta_- \tilde{\varphi}(vt+x)]} :, \quad (17)$$

and the expression of the electric current (5)

$$j(t, x, i) = -\frac{v}{v_F \zeta_+ \sqrt{\pi}} \partial_x \varphi(vt, x, i), \quad (18)$$

needed in the derivation of the conductance. In (16, 17) $: \dots :$ denotes the normal product, whereas

$$\zeta_{\pm} = \sqrt{|\kappa|} \left(\frac{\pi \kappa v_F + 2g_+}{\pi \kappa v_F + 2g_-} \right)^{\pm \frac{1}{4}}, \quad (19)$$

$$v = \frac{\sqrt{(\pi \kappa v_F + 2g_-)(\pi \kappa v_F + 2g_+)}}{\pi |\kappa|}, \quad (20)$$

κ being the *statistical parameter*. The conventional fermionic Luttinger liquid is obtained for $\kappa = 1$. For $\kappa \neq 1$ one has *anyonic* Luttinger liquids. Notice also that the above solution of the Luttinger model is meaningful for coupling constants satisfying $2g_{\pm} > -\pi \kappa v_F$.

C. Symmetry content

The bulk theory, defined by the Lagrangian (2), is invariant under time reversal and global $U(1)$ gauge transformations. Attempting to lift these symmetries to theory on whole star graph Γ , one gets some restrictions on the boundary conditions (7). Time reversal symmetry implies³¹ that $\mathbb{U}$ must be symmetric,

$$\mathbb{U}^t = \mathbb{U}. \quad (21)$$

Concerning the conservation of the $U(1)$ (electric) charge

$$Q = \sum_{i=1}^n \int_0^\infty dx j(t, x, i) \quad (22)$$

on Γ , one knows already from classical electrodynamics that the Kirchhoff's rule

$$\sum_{i=1}^n j(t, 0, i) = 0 \quad (23)$$

must be satisfied at the vertex of Γ . Using (18), one can verify^{20–22} that (23) holds if and only if

$$\sum_{i=1}^n \mathbb{S}_{ij}(k) = 1, \quad \forall j = 1, \dots, n, \quad (24)$$

or, equivalently

$$\sum_{i=1}^n \mathbb{U}_{ij} = 1, \quad \forall j = 1, \dots, n. \quad (25)$$

Combining (13) and (24), one gets

$$\sum_{i=1}^n R_{ij}^{(\eta)} = 0, \quad \forall i\eta \in \mathcal{P}, \quad (26)$$

which will be essential below for checking the Kirchhoff's rule for the conductance away from criticality.

In what follows we assume that both (21) and (25) hold. The case with broken time reversal has been analyzed recently in Ref. 31.

III. BASIC FEATURES OF THE MODEL AWAY FROM CRITICALITY

A. Impact of the analytic structure of $\mathbb{S}(k)$

The analytic properties of the scattering matrix $\mathbb{S}(k)$ deeply influence the physics of the Luttinger liquid on Γ .

The simplest observable, one can investigate in order to illustrate this fact, is the electric current j . More precisely, it is enough to study the relative two-point function, which is fixed²⁹ up to a real parameter τ by causality (local commutativity). Postponing the discussion of

the physical meaning of τ to the end of this subsection, we consider first the current-current correlator following from the definition (18). In the Fock representation of the field φ , defined by equations (A1), (A2) and (A9) in the Appendix A, one finds

$$\langle j(t_1, x_1, i_1) j(t_2, x_2, i_2) \rangle = \frac{v^2}{(2\pi\zeta_+ v_f)^2} [D_{i_1 i_2}(vt_{12}, x_1, x_2) + A_{i_1 i_2}(vt_{12}, x_1, x_2) + B_{i_1 i_2}(vt_1, vt_2, x_1, x_2; \tau)], \quad (27)$$

where

$$D_{i_1 i_2}(t, x_1, x_2) = -\delta_{i_1 i_2} [d^2(t - x_{12}) + d^2(t - x_{12}) + d^2(t + \tilde{x}_{12}) + d^2(t - \tilde{x}_{12})], \quad (28)$$

$$A_{i_1 i_2}(t, x_1, x_2) = \sum_{i\eta \in \mathcal{P}_-} R_{i_1 i_2}^{(\eta)} \{d^2(t + \tilde{x}_{12}) + d^2(t - \tilde{x}_{12}) + \eta [d(t + \tilde{x}_{12}) - d(t - \tilde{x}_{12})] - \eta^2 [w_-(-\eta(t - \tilde{x}_{12}) + w_+(\eta(t + \tilde{x}_{12})))\}, \quad (29)$$

$$B_{i_1 i_2}(t_1, t_2, x_1, x_2; \tau) = \sum_{i\eta \in \mathcal{P}_+} R_{i_1 i_2}^{(\eta)} \{d^2(t_{12} + \tilde{x}_{12}) + d^2(t - \tilde{x}_{12}) + \eta [d(t_{12} + \tilde{x}_{12}) - d(t_{12} - \tilde{x}_{12})] - \eta^2 [w_+(-\eta(t_{12} - \tilde{x}_{12}) + w_- (\eta(t_{12} + \tilde{x}_{12}))) + 2\eta^2 e^{-\eta\tilde{x}_{12}} [\cosh(\eta(\tilde{t}_{12} - 2\tau)) - i \sinh(\eta t_{12})]\}. \quad (30)$$

Here and in what follows $t_{12} = t_1 - t_2$, $x_{12} = x_1 - x_2$, $\tilde{t}_{12} = t_1 + t_2$, $\tilde{x}_{12} = x_1 + x_2$ and

$$d(\xi) = \frac{1}{\xi - i\epsilon}, \quad w_{\pm}(\xi) = e^{-\xi} \text{Ei}(\xi \pm i\epsilon), \quad (31)$$

with $\epsilon > 0$ and Ei the exponential integral function. By construction the functions $A_{i_1 i_2}$ and $B_{i_1 i_2}$ collect the contributions of the antibound and the bound states respectively. The main feature distinguishing these two functions is their time dependence. Notice indeed that $A_{i_1 i_2}$ depends exclusively on t_{12} , whereas $B_{i_1 i_2}$ depends in addition on $\tilde{t}_{12}$ and on the parameter τ . This fact suggest the consideration of two separate cases:

(i) When bound states are absent ($\mathcal{P}_+ = \emptyset$), the behavior of the system is quite orthodox; the function $B_{i_1 i_2}$ vanishes identically and time translations invariance is preserved. Accordingly, the energy is conserved. Because of (21), the theory is also invariant under time reversal

$$t \rightarrow -t. \quad (32)$$

(ii) When bound states are present ($\mathcal{P}_+ \neq \emptyset$) the situation changes drastically. Besides on t_{12} , the dynamics depends also on $\tilde{t}_{12}$, implying that the theory is no longer invariant under time translations. Therefore, the energy is not conserved. The theory depends on the parameter τ as well and is invariant under the time reversal transformation

$$t \rightarrow -t + 2\tau, \quad (33)$$

which is actually a reflection with respect to τ .

The properties collected in point (ii) deserve a more detailed discussion. They imply that for boundary conditions admitting bound states (i.e. some $\alpha_i \in (0, \pi/2)$ - see eqs.(7, 9-12)), the system is not isolated. In fact, there exists a nontrivial energy flow crossing the boundary⁴⁴ at $x = 0$, the direction (outgoing or incoming) being controlled by the parameter τ . In order to illustrate this statement, let us consider the energy density θ of the TL model on Γ . In bosonic coordinates θ takes the form⁴⁵

$$\theta(t, x, i) = \frac{g_-}{\pi\zeta_-^2} (\partial_x \varphi)^2(t, x, i) + \frac{g_+}{\pi\zeta_+^2} (\partial_x \tilde{\varphi})^2(t, x, i). \quad (34)$$

As well known, the field products at coinciding points in (34) contain divergences which must be subtracted. A natural way to fix the subtraction is to take as a reference point the vacuum energy $\langle \theta(t, x) \rangle_{\text{line}}$ on the line.⁴⁶ Using conventional point splitting regularization, one gets in this way²⁹

$$\mathcal{E}(t, x, i) = \langle \theta(t, x, i) \rangle - \langle \theta(t, x) \rangle_{\text{line}} \sim \mathcal{E}(x, i) + \sum_{i\eta \in \mathcal{P}_+} \eta^2 R_{ii}^{(\eta)} e^{-2\eta x} \cosh[2\eta(t - \tau)], \quad (35)$$

where $\mathcal{E}(x, i)$ is a time independent contribution, derived in Ref. 19 but irrelevant for what follows. From (35) we deduce that τ is the instance in which the vacuum energy flow inverts its direction, being outgoing for $t < \tau$ and

incoming for $t > \tau$. This feature is related to the fact that the theory is invariant under the operation (33), which obviously exchanges the two time intervals $(-\infty, \tau)$ and (τ, ∞) . These considerations fix the physical meaning of the parameter τ , which appears in the theory when $\mathcal{P}_+ \neq \emptyset$.

Summarizing, the behavior of off-critical Luttinger junctions is characterized by the two different regimes described in points (i) and (ii) above. In the case (i) one deals with an isolated system in equilibrium. Both energy and electric charge are conserved. The presence of bound states in the regime (ii) significantly modifies this behavior. The energy is no longer conserved because the relative boundary conditions determine a specific boundary interaction with the environment, which drives the system out of equilibrium. A remarkable feature is that the corresponding non-equilibrium state is fixed, up to the value of the parameter τ , by the basic physical requirement of causality.

We stress in conclusion that the above results are obtained in an abstract setup, where we attempt to describe the physics of a quantum wire junction by a Luttinger liquid with specific off-critical boundary conditions at the vertex of the star graph approximating the junction. Further investigations are needed for clarifying the plausibility of these assumptions and the applicability of the results to real-life quantum wire junctions. The case (i) looks more physical since the energy is conserved and bounded from below. In our opinion however, also the regime (ii) can not be excluded a priori, if one considers the possible interactions of the junction with the environment. In order to clarify the situation, one should analyze in the above framework some physical observables, which in principle can be checked experimentally. As an example, we derive in the next subsection the electromagnetic conductance in explicit form. Another attractive possibility to apply the above results is the development of an effective description of the Luttinger liquid on complex quantum wire networks with several junctions and loops, crossed possibly by magnetic fluxes implementing the interaction with the environment. The complete field theory analysis of such networks is usually a complicated problem. One approximate way to face the prob-

lem could be to use the star product approach^{47,48}, the “gluing” technique^{49–51} or transfer matrix formalism⁵² for deriving the *effective* scattering matrix $\mathbb{S}_{\text{eff}}(k)$ relative to the *external* edges of the network. $\mathbb{S}_{\text{eff}}(k)$ admits in general both bound and antibound states and can be used⁵⁰ for constructing a simplified model with one effective off-critical junction.

B. Off-critical conductance

In order to compute the conductance, we couple the Luttinger liquid on Γ to a uniform *classical* electric field⁵³ $E(t, i) = \partial_t A_x(t, i)$, performing in (2) the substitution

$$\partial_x \mapsto \partial_x + iA_x(t, i). \quad (36)$$

This external field coupling deforms the bosonized version of the current according to²⁰

$$j(t, x, i) \mapsto J(t, x, i) = -\frac{v}{v_F \zeta_+ \sqrt{\pi}} \left[\partial_x \varphi(vt, x, i) + \frac{1}{\zeta_+ \sqrt{\pi}} A_x(t, i) \right] \quad (37)$$

The Hamiltonian encoding the interaction of φ and A_x is time dependent and reads²⁰

$$H_{\text{int}}(t) = \frac{1}{\zeta_+ \sqrt{\pi}} \sum_{i=1}^n \int_0^\infty dx (\partial_x \varphi)(vt, x, i) A_x(t, i). \quad (38)$$

We stress that now the system can exchange energy with the environment in two different ways. The first one corresponds to the external force produced by the coupling with the time dependent electric field $E(t, i)$. The second one represents an intrinsic property in the regime (ii), being related to the presence of bound states of $\mathbb{S}(k)$. The conductance, derived below, keeps track of both of them.

In order to derive the conductance, one should compute the expectation value of the current (37) in the external field A_x . This expectation value is given by the following well known⁵⁴ series expansion

$$\begin{aligned} \langle J(t, x, i) \rangle_{A_x} &= \langle J(t, x, i) \rangle - i \int_{-\infty}^t d\tau_1 \langle [J(t, x, i), H_{\text{int}}(\tau_1)] \rangle + \cdots \\ &+ (-i)^n \int_{-\infty}^t d\tau_1 \int_{-\infty}^{\tau_1} d\tau_2 \cdots \int_{-\infty}^{\tau_{(n-1)}} d\tau_n [\cdots [[J(t, x, i), H_{\text{int}}(\tau_1)], H_{\text{int}}(\tau_2)], \cdots, H_{\text{int}}(\tau_n)] + \cdots \end{aligned} \quad (39)$$

in powers of H_{int} . This expansion is particularly simple in our case. In fact, from (37,38) one obtains that

$$[[J(t, x, i), H_{\text{int}}(\tau_1)], H_{\text{int}}(\tau_2)] = 0, \quad (40)$$

cutting the infinite series (39) to the sum of the first two terms, namely

$$\langle J(t, x, i) \rangle_{A_x} = \frac{v}{v_F \zeta_+^2 \pi} \left[A_x(t, i) + i \sum_{j=1}^n \int_{-\infty}^t d\tau \int_0^\infty dy A_y(\tau, j) \langle [\partial_y \varphi(v\tau, y, j), \partial_x \varphi(vt, x, i)] \rangle \right]. \quad (41)$$

The expectation value $\langle J(t, x, i) \rangle_{A_x}$ is therefore linear in A_x , implying that the linear response approximation⁵⁴ to $\langle J(t, x, i) \rangle_{A_x}$ is actually exact in this case.

Let us suppose now that the external field is switched on at $t = t_0$, i.e. $A_x(t, i) = 0$ for $t < t_0$. Using the representation (A15) for the commutator in the left hand side of (41), one finds for $t > t_0$

$$\langle J(t, 0, i) \rangle_{A_x} = G_{\text{line}} \sum_{j=1}^n \int_{-\infty}^\infty \frac{d\omega}{2\pi} \hat{A}_x(\omega, j) e^{-i\omega t} \left[\delta_{ji} - S_{ji} \left(\frac{\omega}{v} \right) - \sum_{i\eta \in \mathcal{P}} R_{ji}^{(\eta)} \frac{v\eta}{v\eta + i\omega} e^{(t-t_0)(v\eta + i\omega)} \right], \quad (42)$$

where

$$G_{\text{line}} = \frac{v}{v_F \zeta_+^2 \pi}, \quad \hat{A}_x(\omega, i) = \int_{-\infty}^\infty d\tau e^{i\omega\tau} A_x(\tau, i). \quad (43)$$

The conductance can be extracted directly from (42). The result is

$$G_{ij}(\omega, t - t_0) = G_{\text{line}} \left[\delta_{ij} - S_{ij} \left(\frac{\omega}{v} \right) - \sum_{i\eta \in \mathcal{P}} R_{ij}^{(\eta)} \frac{v\eta}{v\eta + i\omega} e^{(t-t_0)(v\eta + i\omega)} \right], \quad t > t_0, \quad (44)$$

which represents the off-critical generalization of the formula (1) we are looking for. Notice that away from criticality $G_{ij}(\omega, t - t_0)$ is in general complex, showing that off-critical junctions may have a nontrivial impedance. Since eq. (41) involves the commutator of the field φ , the conductance does not depend on the parameter τ , but depends on the time $t - t_0$ elapsed after switching on the external field.

Before discussing the main features of (44), we would like to mention two useful checks. First of all we observe that at criticality, where scale invariance implies that $\mathbb{S}$ is constant (k -independent), the sum over the poles vanishes and (44) precisely reproduces the expression (1) from the introduction. A second highly nontrivial check is the Kirchhoff rule

$$\sum_{i=1}^n G_{ij}(\omega, t - t_0) = 0, \quad (45)$$

which is a consequence of the conservation of the electric charge, namely of equations (24, 26).

The sum in the right hand side of (44) runs over both negative and positive poles $\mathcal{P} = \mathcal{P}_- \cup \mathcal{P}_+$. The antibound states $\mathcal{P}_-$ produce damped oscillations in $t - t_0$. If bound states are absent ($\mathcal{P}_+ = \emptyset$),

$$\lim_{t \rightarrow \infty} G_{ij}(\omega, t - t_0) = \lim_{t_0 \rightarrow -\infty} G_{ij}(\omega, t - t_0) = G_{\text{line}} \left[\delta_{ij} - S_{ij} \left(\frac{\omega}{v} \right) \right], \quad (46)$$

which gives the conductance one will observe in the regime (i) long time after switching on the external field.

Finally, the bound states $\mathcal{P}_+$ give origin to oscillations whose amplitude is growing exponentially with $t - t_0$. The oscillations in (44) provide therefore a nice experimental signature for testing the analytic structure of the scattering matrix $\mathbb{S}(k)$.

IV. OUTLOOK AND CONCLUSIONS

Off-critical Luttinger junctions are characterized by a scattering matrix $\mathbb{S}(k)$ which admits in general bound and antibound states. The presence/absence of bound states determines two different regimes of the theory. A scattering matrix without bound states gives rise to an isolated equilibrium system. One bound state is enough to change radically the situation. We have shown in fact that each such a state generates an oscillator degree of freedom, whose contribution is fixed by local commutativity up to a common free parameter τ . These additional degrees of freedom break the invariance under time translations and drive the system out of equilibrium. Accordingly, the vacuum energy is time dependent: it decays exponentially in the interval $(-\infty, \tau)$ and grows at the same rate in (τ, ∞) . These two intervals are related by time reversal and are characterized by a nontrivial outgoing and incoming vacuum energy flows. Recalling that the traditional way⁵⁵ for driving a system out of equilibrium is to couple it with a “bath” of external oscillators, we have shown above that under certain conditions such oscillators can be automatically generated by boundary effects. Junctions with bound states provide therefore

an intrinsic mechanism for constructing non-equilibrium quantum systems, whose behavior is governed by purely boundary phenomena. Since this mechanism is based on the general physical requirement of causality, it extends⁵⁶ also to systems in space dimensions greater than one.

The off-critical conductance formula (44) has various interesting properties. A first remarkable feature of (44) is the different impact of antibound and bound states, which hopefully can be tested experimentally. The result (44) turns out to be useful also at the level of effective theory for quantum wire networks with several junctions, where it applies for the effective scattering matrix $\mathbb{S}_{\text{eff}}(k)$ associated to the external edges of the network.

Acknowledgments

We thank Pasquale Calabrese, Benoit Douçot, Inès Saffi and Robert Schrader for enlightening discussions and correspondence. The research of B. B. has been supported in part by the NSF grant PHY-0757868.

Appendix A: Effect of the bound states of $\mathbb{S}(k)$ on the field φ

We assume below that $\mathbb{S}(k)$ admits bound states, the case $\mathcal{P}_+ = \emptyset$ being standard¹⁹. When $\mathcal{P}_+ \neq \emptyset$ it is natural to represent the field φ as a linear combination

$$\varphi(t, x, i) = \varphi^{(s)}(t, x, i) + \varphi^{(b)}(t, x, i), \quad (\text{A1})$$

where $\varphi^{(s)}$ collects the contribution of the *scattering* states of $\mathbb{S}(k)$ and $\varphi^{(b)}$, that of its *bound* states. The scattering component $\varphi^{(s)}$ is known from previous studies¹⁹. One has

$$\varphi^{(s)}(t, x, i) = \int_{-\infty}^{\infty} \frac{dk}{2\pi\sqrt{2|k|}} \left[a_i^*(k) e^{i(|k|t - kx)} + a_i(k) e^{-i(|k|t - kx)} \right], \quad (\text{A2})$$

where $\{a_i(k), a_i^*(k) : k \in \mathbb{R}\}$ generate the reflection-transmission algebra^{40–42} $\mathcal{A}$

$$\begin{aligned} [a_i(k), a_j(p)] &= [a_i^*(k), a_j^*(p)] = 0, \\ [a_i(k), a_j^*(p)] &= 2\pi[\delta(k-p)\delta_{ij} + \mathbb{S}_{ij}(k)\delta(k+p)], \end{aligned} \quad (\text{A3})$$

and satisfy the constraints

$$a_i(k) = \sum_{j=1}^n \mathbb{S}_{ij}(k) a_j(-k), \quad (\text{A4})$$

$$a_i^*(k) = \sum_{j=1}^n a_j^*(-k) \mathbb{S}_{ji}(-k), \quad (\text{A5})$$

which are consistent because unitarity and Hermitian analyticity of $\mathbb{S}(k)$ imply $\mathbb{S}(k)\mathbb{S}(-k) = \mathbb{I}$.

Since for $\mathcal{P}_+ \neq \emptyset$ the scattering states are not complete, the scattering component $\varphi^{(s)}$ is neither a canonical nor a local field. In fact, keeping in mind that $\tilde{x}_{12} > 0$ and using the Cauchy integral formula one finds

$$\begin{aligned} [(\partial_t \varphi^{(s)})(0, x_1, i_1), \varphi^{(s)}(0, x_2, i_2)] &= \\ -i\delta_{i_1 i_2} \delta(x_1 - x_2) + i \sum_{i\eta \in \mathcal{P}_+} \eta e^{-\eta \tilde{x}_{12}} R_{i_1 i_2}^{(\eta)}, \end{aligned} \quad (\text{A6})$$

and, at space-like separations $t_{12}^2 - x_{12}^2 < 0$,

$$\begin{aligned} [\varphi^{(s)}(t_1, x_1, i_1), \varphi^{(s)}(t_2, x_2, i_2)] &= \\ i \sum_{i\eta \in \mathcal{P}_+} e^{-\eta \tilde{x}_{12}} \sinh(\eta t_{12}) R_{i_1 i_2}^{(\eta)}. \end{aligned} \quad (\text{A7})$$

We turn now to the construction of the boundary component $\varphi^{(b)}$ which is designed in such a way that the total field (A1) is both canonical and local. Let us introduce first the set $\mathcal{I}_+ = \{i : \eta_i > 0\}$ labeling the *different* poles of $\mathbb{S}(k)$ in the upper half plane. Then the normalizable solutions of the wave equation on Γ corresponding to the bound states are $\{e^{-\eta_i(x \pm t)} : i \in \mathcal{I}_+\}$. The main idea now is to associate with each index $i \in \mathcal{I}_+$ a quantum oscillator. The quantum boundary degrees of freedom are described therefore by the algebra $\mathcal{B}$ generated by $\{b_i, b_i^* : i \in \mathcal{I}_+\}$, which satisfy

$$[b_{i_1}, b_{i_2}] = [b_{i_1}^*, b_{i_2}^*] = 0, \quad [b_{i_1}, b_{i_2}^*] = \delta_{i_1 i_2}, \quad (\text{A8})$$

and commute with $\{a_i(k), a_i^*(k)\}$. Now, the field $\varphi^{(b)}$ is defined by²⁹

$$\begin{aligned} \varphi^{(b)}(t, x, i) &= \frac{1}{\sqrt{2}} \sum_{j \in \mathcal{I}_+} \mathcal{U}_{ij} [(b_j^* + b_j) e^{-\eta_j(x+t-\tau)} + \\ &\quad i(b_j^* - b_j) e^{-\eta_j(x-t+\tau)}], \end{aligned} \quad (\text{A9})$$

$\mathcal{U}$ being the unitary matrix which appears in (9). By construction the field (A9) satisfies the equation of motion (6), the boundary condition (7) and depends on the parameter $\tau \in \mathbb{R}$, whose physical meaning has been discussed in section II.D. One can also verify²⁹ that the total field (A1) is both canonical and local. In fact,

$$\begin{aligned} [\varphi^{(b)}(t_1, x_1, i_1), \varphi^{(b)}(t_2, x_2, i_2)] &= \\ -i \sum_{i\eta \in \mathcal{P}_+} e^{-\eta \tilde{x}_{12}} \sinh(\eta t_{12}) R_{i_1 i_2}^{(\eta)}, \end{aligned} \quad (\text{A10})$$

which precisely compensates the right hand side of (A7).

The Hamiltonian H , generating the time evolution of φ according to

$$[H, \varphi(t, x, i)] = -i(\partial_t \varphi)(t, x, i). \quad (\text{A11})$$

is given by

$$H = H^{(s)} + H^{(b)}, \quad (\text{A12})$$

where the scattering contribution $H^{(s)}$ has the standard form

$$H^{(s)} = \frac{1}{2} \sum_{i=1}^n \int_{-\infty}^{\infty} \frac{dk}{2\pi} |k| a_i^*(k) a_i(k), \quad (\text{A13})$$

and

$$H^{(b)} = \frac{i}{2} \sum_{j \in \mathcal{I}_+} \eta_j (b_j^2 - b_j^{*2}). \quad (\text{A14})$$

We see that $H^{(b)}$ is not at all the Hamiltonian of the conventional harmonic oscillator, which is the origin of the peculiar time dependence of the vacuum energy density (35).

For deriving the conductance it is convenient to plug in (41) the following integral representation of the commutator

$$[\varphi(t_1, x_1, i_1), \varphi(t_2, x_2, i_2)] = -i \int_{-\infty}^{\infty} \frac{dk}{2\pi} \frac{1}{k} \sin(kt_{12}) \left[e^{ikx_{12}} \delta_{i_1 i_2} + e^{ik\tilde{x}_{12}} S_{i_1 i_2}(k) \right] - i \sum_{i\eta \in \mathcal{P}_+} e^{-\eta \tilde{x}_{12}} \sinh(\eta t_{12}) R_{i_1 i_2}^{(\eta)}. \quad (\text{A15})$$

We stress that (A15) is τ -independent. Finally, the current-current correlator (27-31) is obtained in the Fock representation of the algebras $\mathcal{A}$ and $\mathcal{B}$, characterized as usual by a vacuum state Ω annihilated by $a_i(k)$ and b_i .

In conclusion, we would like to stress once more the

fundamental role played by causality (local commutativity) in the above construction. In fact, this deep physical requirement fixes (up to the parameter τ) the whole structure of the field φ and its time evolution.

-
- ¹ C. L. Kane and M. P. A. Fisher, Phys. Rev. Lett. **68**, 1220 (1992); Phys. Rev. B **46**, 15233 (1992).
 - ² I. Safi, H. J. Schulz, Phys. Rev. B **52**, R17040 (1995).
 - ³ C. Nayak, M. P. A. Fisher, A. W. W. Ludwig and H. H. Lin, Phys. Rev. B **59**, 15694 (1999).
 - ⁴ I. Safi, P. Devillard, and T. Martin, Phys. Rev. Lett. **86**, 4628 (2001).
 - ⁵ J.E. Moore and X.-G. Wen, Phys. Rev. B **66**, 115305 (2002).
 - ⁶ H. Yi, Phys. Rev. B **65**, 195101 (2002).
 - ⁷ S. Lal, S. Rao, and D. Sen, Phys. Rev. B **66**, 165327 (2002).
 - ⁸ S. Chen, B. Trauzettel, and R. Egger, Phys. Rev. Lett. **89**, 226404 (2002).
 - ⁹ K-V. Pham, F. Piechon, K-I Imura, P. Lederer, Phys. Rev. B **68**, 205110 (2003).
 - ¹⁰ C. Chamon, M. Oshikawa, and I. Affleck, Phys. Rev. Lett. **91**, 206403 (2003); M. Oshikawa, C. Chamon, and I. Affleck, J. Stat. Mech. P02008 (2006).
 - ¹¹ F. Dolcini, H. Grabert, I. safi and B. Trauzettel, Phys. Rev. Lett. **91**, 266402 (2003).
 - ¹² S. Rao and D. Sen, Phys. Rev. B **70**, 195115 (2004).
 - ¹³ K. Kazymyrenko and B. Douçot, Phys. Rev. B **71**, 075110 (2005).
 - ¹⁴ E.-A. Kim, M. J. Lawler, S. Vishveshwara, E. Fradkin, Phys. Rev. Lett. **95**, 176402 (2005); Phys. Rev. B **74**, 155324 (2006).
 - ¹⁵ D. Giuliano and P. Sodano, Nucl. Phys. B **711**, 480 (2005).
 - ¹⁶ T. Enss, V. Meden, S. Andergassen, X. Barnabe-Theriat, W. Metzner, K. Schonhammer, Phys. Rev. B **71**, 155401 (2005); X. Barnabe-Theriat, A. Sedeki, V. Meden, K. Schonhammer, Phys. Rev. Lett. **94**, 136405 (2005); X. Barnabe-Theriat, A. Sedeki, V. Meden, K. Schonhammer, Phys. Rev. B **71**, 205327 (2005).
 - ¹⁷ D. Friedan, cond-mat/0505084; cond-mat/0505085.
 - ¹⁸ S. Das, S. Rao, D. Sen, Phys. Rev. B **74**, 045322 (2006).
 - ¹⁹ B. Bellazzini and M. Mintchev, J. Phys. A **39**, 11101 (2006) [arXiv:hep-th/0605036].
 - ²⁰ B. Bellazzini, M. Mintchev and P. Sorba, J. Phys. A **40**, 2485 (2007) [arXiv:hep-th/0611090].
 - ²¹ B. Bellazzini, M. Burrello, M. Mintchev and P. Sorba, Proc. Symp. Pure Math. **77**, 639 (2008), arXiv:0801.2852 [hep-th].
 - ²² B. Bellazzini, P. Calabrese and M. Mintchev, Phys. Rev. B **79** 085122 (2009), [arXiv:0808.2719].
 - ²³ C.-Y. Hou and C. Chamon, Phys. Rev. B **77**, 155422 (2008).
 - ²⁴ S. Das, S. Rao, and A. Saha, Phys. Rev. B **77**, 155418 (2008); Europhys. Lett. **81**, 67001 (2008).
 - ²⁵ S. Das and S. Rao, Phys. Rev. B **78**, 205421 (2008).
 - ²⁶ C.-Y. Hou, E.-A. Kim, and C. Chamon, Phys. Rev. Lett. **102**, 076602 (2009).
 - ²⁷ A. Agarwal, S. Das, S. Rao and D. Sen, Phys. Rev. Lett. **103**, 026401 (2009); Erratum, Phys. Rev. Lett. **103**, 079903 (2009), arXiv:0810.3513 [cond-mat].
 - ²⁸ S. Das, S. Rao and A. Saha, Phys. Rev. B **79**, 155416 (2009).
 - ²⁹ B. Bellazzini, M. Mintchev and P. Sorba, J. Math. Phys. (in press), arXiv:0810.3101 [hep-th].
 - ³⁰ I. Safi, arXiv:0906.2363 [cond-mat].
 - ³¹ B. Bellazzini, M. Mintchev and P. Sorba, Phys. Rev. B **80**, 25441 (2009), arXiv:0907.4221 [hep-th].
 - ³² A. Rahmani, C.-Y. Hou, C. Chamon and I. Affleck, arXiv:1001.3877 [cond-mat].
 - ³³ F. D. M. Haldane, Phys. Rev. Lett. **47**, 1840 (1981); J. Phys. C **15**, 2585 (1981). See for instance also A. O. Gogolin, A. A. Nersisyan, and A. M. Tsvelik, Bosonization and strongly correlated systems (Cambridge University Press, Cambridge, U.K., 2004).

- ³⁴ Sometimes the combinations $g_{2,4} = 2(g_- \mp g_+)$ are used as well.
- ³⁵ V. Kostykin and R. Schrader, Fortschr. Phys. **48**, 703 (2000).
- ³⁶ P. Kuchment, arXiv:0802.3442 [math-ph].
- ³⁷ M. Harmer, J. Phys. A **33** (2000) 9015.
- ³⁸ The boundary condition (7) can be reformulated in terms of currents also away from criticality.²⁹
- ³⁹ A. Liguori, M. Mintchev and L. Zhao, Commun. Math. Phys. **194**, 569 (1998) [arXiv:hep-th/9607085].
- ⁴⁰ M. Mintchev, E. Ragoucy and P. Sorba, Phys. Lett. B **547**, 313 (2002) [arXiv:hep-th/0209052].
- ⁴¹ M. Mintchev, E. Ragoucy and P. Sorba, J. Phys. A **36**, 10407 (2003) [arXiv:hep-th/0303187].
- ⁴² M. Mintchev and P. Sorba, J. Stat. Mech. **0407**, P001 (2004) [arXiv:hep-th/0405264].
- ⁴³ Because of their origin, these states are called *boundary* (anti)bound states as well.
- ⁴⁴ We stress that this is a purely boundary effect and that the energy momentum tensor in the bulk of Γ satisfies the continuity equation.
- ⁴⁵ G. F. Dell-Antonio, Y. Frishman and D. Zwanziger, Phys. Rev. D **6**, 988 (1972).
- ⁴⁶ This is a standard procedure¹⁹ for deriving the Casimir energy density on the graph Γ .
- ⁴⁷ V. Kostykin and R. Schrader, J. Math. Phys. **42** (2001) 1563.
- ⁴⁸ R. Schrader, J. Phys. A **42** (2009) 495401, [arXiv:0907.1522 [hep-th]].
- ⁴⁹ M. Mintchev and E. Ragoucy, J. Phys. A **40** (2007) 9515 [arXiv:0705.1322 [hep-th]].
- ⁵⁰ E. Ragoucy, J. Phys. A **42** (2009) 295205, [arXiv:0901.2431 [hep-th]].
- ⁵¹ V. Caudrelier and E. Ragoucy, Nucl. Phys. B **828** (2010) 515, [arXiv:0907.5359 [math-ph]].
- ⁵² S. Khachatryan, A. Sedrakyan and P. Sorba, Nucl. Phys. B **825** (2010) 444, [arXiv:0904.2688 [cond-mat.mes-hall]].
- ⁵³ We use the Weyl gauge $A_t = 0$.
- ⁵⁴ A. L. Fetter and J. D. Walecka, *Quantum theory of many-particle systems*, (Dover Publications, New York, 2003).
- ⁵⁵ A. O. Caldeira and A. J. Legget, Annals Phys. **149** (1983) 374.
- ⁵⁶ B. Bellazzini, M. Mintchev and P. Sorba, (work in progress).